


Meclizine HCl Tablets, USP Rx Only

DESCRIPTION

Chemically, Meclizine HCl is 1-(*p*-chloro- α -phenylbenzyl)-4-(*m*-methylbenzyl) piperazine dihydrochloride monohydrate.



$C_{25}H_{27}ClN_2 \cdot 2HCl \cdot H_2O$

Molecular Weight 481.88

Meclizine HCl Tablets, USP are available in two different strengths: 12.5 mg and 25 mg. In addition each tablet contains the following inactive ingredients: Colloidal Silicon Dioxide, Croscarmellose Sodium, Lactose Monohydrate, Magnesium Stearate, Microcrystalline Cellulose. Also, Meclizine HCl Tablets USP, 12.5 mg contains FD&C Blue #1 Aluminum Lake (11-13%) and Meclizine HCl Tablets USP, 25 mg contains D&C Yellow #10 Aluminum Lake (15-20%).

CLINICAL PHARMACOLOGY

Meclizine HCl is an antihistamine which shows marked protective activity against nebulized histamine and lethal doses of intravenously injected histamine in guinea pigs. It has a marked effect in blocking the vasodepressor response to histamine, but only a slight blocking action against acetylcholine. Its activity is relatively weak in inhibiting the spasmogenic action of histamine on isolated guinea pig ileum.

INDICATIONS AND USAGE

For the management of nausea and vomiting, and dizziness associated with motion sickness.

CONTRAINDICATIONS

Meclizine HCl is contraindicated in individuals who have shown a previous hypersensitivity to it.

WARNINGS

Since drowsiness may, on occasion, occur with use of this drug, patients should be warned of this possibility and cautioned against driving a car or operating dangerous machinery.

Patients should avoid alcoholic beverages while taking this drug.

Due to its potential anticholinergic action, this drug should be used with caution in patients with asthma, glaucoma or enlargement of the prostate gland.

PRECAUTIONS

PREGNANCY, Teratogenic Effects

Pregnancy Category B. Reproduction studies in rats have shown cleft palates at 25-50 times the human dose. Epidemiological studies in pregnant women, however, do not indicate that medicine increases the risk of abnormalities when administered during pregnancy. Despite the animal findings, it would appear that the possibility of fetal harm is remote. Nevertheless, meclizine, or any other medication, should be used during pregnancy only if clearly necessary.

Pediatric Use

Clinical studies establishing safety and effectiveness in children have not been done; therefore, usage is not recommended in children under 12 years of age.

ADVERSE REACTIONS

Drowsiness, dry mouth and, on rare occasions, blurred vision have been reported.

DOSAGE AND ADMINISTRATION

Motion Sickness

The initial dose of 25 to 50 mg of Meclizine HCl should be taken one hour prior to travel for protection against motion sickness. Thereafter, the dose may be repeated every 24 hours for the duration of the journey.

How Supplied

Meclizine HCl Tablets, USP are available in the following strengths and package sizes:

| | |
|--|------------------|
| 12.5 mg (Blue, oval-shaped, scored, debossed with TL122) | |
| Bottles of 100 | NDC 59746-122-06 |
| Bottles of 1000 | NDC 59746-122-10 |

| | |
|--|------------------|
| 25 mg (Yellow, oval-shaped, scored, debossed with TL121) | |
| Bottles of 100 | NDC 59746-121-06 |
| Bottles of 1000 | NDC 59746-121-10 |

Store at 20-25°C (68-77°F) (See USP Controlled Room Temperature).

Manufactured By:

Jubilant Cadista Pharmaceuticals Inc.
Salisbury, MD 21801, USA

Revised 03/11

