

# PANTOPRAZOLE SODIUM DELAYED-RELEASE TABLETS USP

(pan toe' pra zole soe' dee um)

## What is the most important information I should know about pantoprazole sodium delayed-release tablets?

You should take pantoprazole sodium delayed-release tablets exactly as prescribed, at the lowest dose possible and for the shortest time needed.

Pantoprazole sodium delayed-release tablets may help your acid-related symptoms, but you could still have serious stomach problems. Talk with your doctor.

## Pantoprazole sodium delayed-release tablets can cause serious side effects, including:

- **A type of kidney problem (acute interstitial nephritis).** Some people who take proton pump inhibitor (PPI) medicines, including pantoprazole sodium delayed-release tablets, may develop a kidney problem called acute interstitial nephritis that can happen at any time during treatment with pantoprazole sodium delayed-release tablets. Call your doctor right away if you have a decrease in the amount that you urinate or if you have blood in your urine.
- **Diarrhea caused by an infection (*Clostridium difficile*) in your intestines.** Call your doctor right away if you have watery stools or stomach pain that does not go away. You may or may not have a fever.
- **Bone fractures (hip, wrist, or spine).** Bone fractures in the hip, wrist, or spine may happen in people who take multiple daily doses of PPI medicines and for a long period of time (a year or longer). Tell your doctor if you have a bone fracture, especially in the hip, wrist, or spine.
- **Certain types of lupus erythematosus.** Lupus erythematosus is an autoimmune disorder (the body's immune cells attack other cells or organs in the body). Some people who take PPI medicines, including pantoprazole sodium delayed-release tablets, may develop certain types of lupus erythematosus or have worsening of the lupus they already have. Call your doctor right away if you have new or worsening joint pain or a rash on your cheeks or arms that gets worse in the sun.

Talk to your doctor about your risk of these serious side effects. Pantoprazole sodium delayed-release tablets can have other serious side effects. See **“What are the possible side effects of pantoprazole sodium delayed-release tablets?”**

## What are pantoprazole sodium delayed-release tablets?

A prescription medicine called a proton pump inhibitor (PPI) used to reduce the amount of acid in your stomach.

**In adults**, pantoprazole sodium delayed-release tablets are used for:

- up to 8 weeks for the healing and symptom relief of acid-related damage to the lining of the esophagus (called erosive esophagitis or EE). Your doctor may prescribe another 8 weeks of pantoprazole sodium delayed-release tablets in patients whose EE does not heal.
- maintaining healing of EE and to help prevent the return of heartburn symptoms caused by GERD. It is not known if pantoprazole sodium delayed-release tablets are safe and effective when used for longer than 12 months for this purpose.
- the long-term treatment of conditions where your stomach makes too much acid. This includes a rare condition called Zollinger-Ellison Syndrome.

**In children 5 years of age and older**, pantoprazole sodium delayed-release tablets are used for:

- up to 8 weeks for the healing and symptom relief of EE. It is not known if pantoprazole sodium delayed-release tablets are safe if used longer than 8 weeks in children.

Pantoprazole sodium delayed-release tablets are not for use in children under 5 years of age.

It is not known if pantoprazole sodium delayed-release tablets are safe and effective in children for treatment other than EE.

## Do not take pantoprazole sodium delayed-release tablets if you are:

- allergic to pantoprazole sodium, any other PPI medicine, or any of the ingredients in pantoprazole sodium delayed-release tablets. See the end of this Medication Guide for a complete list of ingredients.
- taking a medicine that contains rilpivirine (EDURANT, COMPLERA, ODEFSEY) used to treat HIV-1 (Human Immunodeficiency Virus).

## Before taking pantoprazole sodium delayed-release tablets, tell your doctor about all of your medical conditions, including if you:

- have low magnesium levels in your blood.
- are pregnant or plan to become pregnant. Pantoprazole sodium delayed-release tablets may harm your unborn baby. Tell your doctor if you become pregnant or think you may be pregnant during treatment with pantoprazole sodium delayed-release tablets.
- are breastfeeding or plan to breastfeed. Pantoprazole can pass into your breast milk. You and your doctor should decide if you will take pantoprazole sodium delayed-release tablets or breastfeed. You should not do both. Talk with your doctor about the best way to feed your baby if you take pantoprazole sodium delayed-release tablets.

Item Code: 50000002915		Superseded Item Code: 50000002806		Pharma Code: NA	
Component: Medication Guide		Substrate: 40 GSM Bible Paper			
Dimension	Foil width: NA		Blister (LxW): NA		
	Print Registration: NA				
	Carton (LxWxH): NA				
	PIL/ Medication Guide		Open (LxW): 280 x 216 mm		
			Folded (LxW): NA		
Label (LxW): NA					
Client & Country : Cadista- US		Market: US		Reason for Artwork: RLD UPDATE 20th Dec, 2017	
Reference Spec No: PS2861		Pantone: <span style="display:inline-block; width:10px; height:10px; background-color:black;"></span> Black <span style="display:inline-block; width:10px; height:10px; background-color:magenta;"></span> Dieline			
Special Instruction (If any): NA					
Site Packaging Development Sign and Date		Production Sign and Date		QA Sign and Date	



**Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins and herbal supplements. **Especially tell your doctor if you take methotrexate (Otrexup, Rasuvo, Trexall, XATMEP), digoxin (LANOXIN), or a water pill (diuretic).**

**How should I take pantoprazole sodium delayed-release tablets?**

- Take pantoprazole sodium delayed-release tablets exactly as prescribed by your doctor.
  - **Do not split, chew, or crush pantoprazole sodium delayed-release tablets.**
  - Swallow pantoprazole sodium delayed-release tablet whole, with or without food.
  - Tell your doctor if you are not able to swallow your pantoprazole sodium delayed-release tablet.
  - You may use antacids while taking pantoprazole sodium delayed-release tablets.
- If you miss a dose of pantoprazole sodium delayed-release tablets, take it as soon as possible. If it is almost time for your next dose, do not take the missed dose. Take the next dose at your regular time. Do not take 2 doses at the same time.
- If you take too much pantoprazole sodium delayed-release tablets, call your doctor or your poison control center at 1-800-222-1222 right away or go to the nearest emergency room.

**What are the possible side effects of pantoprazole sodium delayed-release tablets?**

**Pantoprazole sodium delayed-release tablets can cause serious side effects, including:**

- **See “What is the most important information I should know about pantoprazole sodium delayed-release tablets?”**
- **Low vitamin B-12 levels** in your body can happen in people who have taken pantoprazole sodium delayed-release tablets for a long time (more than 3 years). Tell your doctor if you have symptoms of low vitamin B-12 levels, including shortness of breath, lightheadedness, irregular heartbeat, muscle weakness, pale skin, feeling tired, mood changes, and tingling or numbness in the arms and legs.
- **Low magnesium levels in your body** can happen in people who have taken pantoprazole sodium delayed-release tablets for at least 3 months. Tell your doctor if you have symptoms of low magnesium levels, including seizures, dizziness, irregular heartbeat, jitteriness, muscle aches or weakness, and spasms of hands, feet or voice.

**The most common side effects of pantoprazole sodium delayed-release tablets in adults include:** headache, diarrhea, nausea, stomach-area (abdominal) pain, vomiting, gas, dizziness, and joint pain.

**The most common side effects of pantoprazole sodium delayed-release tablets in children include:** upper respiratory infection, headache, fever, diarrhea, vomiting, rash, and stomach-area (abdominal) pain.

These are not all the possible side effects of pantoprazole sodium delayed-release tablets. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store pantoprazole sodium delayed-release tablets?**

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].

**Keep pantoprazole sodium delayed-release tablets and all medicines out of the reach of children.**

**General information about the safe and effective use of pantoprazole sodium delayed-release tablets.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use pantoprazole sodium delayed-release tablets for a condition for which it was not prescribed. Do not give pantoprazole sodium delayed-release tablets to other people, even if they have the same symptoms that you have. It may harm them. You can ask your doctor or pharmacist for information about pantoprazole sodium delayed-release tablets that is written for health professionals.

For more information, call toll-free 1-800-313-4623.

**What are the ingredients in pantoprazole sodium delayed-release tablets?**

**Active ingredient:** pantoprazole sodium sesquihydrate USP

**Inactive ingredients:** calcium stearate, ferric oxide red, ferric oxide black, hypromellose, hypromellose phthalate, mannitol, methacrylic acid copolymer, polyethylene glycol, silicon dioxide, sodium carbonate, sucrose, talc, titanium dioxide, and triethyl citrate. In addition to the above, printing ink used to imprint the delayed-release tablet contains ammonium hydroxide, ferric oxide black, propylene glycol, and shellac.

**This Medication Guide has been approved by the U.S. Food and Drug Administration.**

All brand names listed are the registered trademarks of their respective owners and are not trademarks of Jubilant Generics Limited.

**Rx Only**

**Manufactured by:**  
Jubilant Generics Limited  
Roorkee - 247661, India

**Marketed by:**  
Jubilant Cadista Pharmaceuticals Inc.  
Salisbury, MD 21801, USA

**Revised: 02/2018**

50000002915

Neutral Code No. UK/Drugs/58/UA/2007

Item Code: 50000002915		Superseded Item Code: 50000002806		Pharma Code: NA	
Component: Medication Guide		Substrate: 40 GSM Bible Paper			
Dimension	Foil width: NA		Blister (LxW): NA		
	Print Registration: NA				
	Carton (LxWxH): NA				
	PIL/ Medication Guide		Open (LxW): 280 x 216 mm		
			Folded (LxW): NA		
Label (LxW): NA					
Client & Country : Cadista- US		Market: US		Reason for Artwork: RLD UPDATE 20th Dec, 2017	
Reference Spec No: PS2861		Pantone: <span style="display: inline-block; width: 10px; height: 10px; background-color: black;"></span> Black <span style="display: inline-block; width: 10px; height: 10px; background-color: magenta;"></span> Dieline			
Special Instruction (If any): NA					
Site Packaging Development Sign and Date		Production Sign and Date		QA Sign and Date	